

CENTRE DE PHYSIQUE THEORIQUE - CNRS - Luminy, Case 907

F-13288 Marseille Cedex 9 - France

Unité Propre de Recherche 7061

The angular momentum operator in the Dirac equation

Víctor M. Villalba¹

Abstract

The Dirac equation in spherically symmetric fields is separated in two different tetrad frames. One is the standard cartesian (fixed) frame and the second one is the diagonal (rotating) frame. After separating variables in the Dirac equation in spherical coordinates, and solving the corresponding eigenvalues equations associated with the angular operators, we obtain that the spinor solution in the rotating frame can be expressed in terms of Jacobi polynomials, and it is related to the standard spherical harmonics, which are the basis solution of the angular momentum in the Cartesian tetrad, by a similarity transformation

Key-Words : 3.65, 11.10, 4.90.

April 1994

CPT-94/P.3027

anonymous ftp or gopher: cpt.univ-mrs.fr

¹permanent address: Centro de Física, Instituto Venezolano de Investigaciones Científicas IVIC, Apdo. 21827, Caracas 1020-A Venezuela
e-mail address: villalba@cpt.univ-mrs.fr, villalba@dino.conicit.ve

1 Introduction

The Dirac equation is a system of four coupled partial differential equation which describes the relativistic electron and other spin 1/2 particles. Despite the remarkable effort made during the last decades in order to find exact solutions for the relativistic electron in the presence of external fields, the amount of solvable problems is relatively scarce, being the Coulomb problem [1], the plane wave [2], and some electromagnetic configurations [3] the most representative examples. The problem of finding exact solutions of the Dirac equation is closely related to the possibility of accomplishing a complete separation of variables of the Dirac equation.

During the last decades there have been different approaches to tackling the problem of separating variables in the Dirac equation in the presence of external fields; we have the Stäckel spaces method developed by Bagrov and co-workers ([3] based on the experienced acquired in separating variables in the Klein Gordon equation), the theory of R-separability by Miller [4, 5], and matrix modification of the Stäckel proposed by Cook [6] Recently an algebraic method of separation of variables [9, 10] has been successfully applied in different gravitational and electromagnetic configurations. The main idea of the algebraic method of separation is to try to reduce the original Dirac equation, which in the presence of electromagnetic fields takes the form

$$\{D\}\Psi = \{\tilde{\gamma}^\mu(\partial_\mu - \Gamma_\mu - iA_\mu) + m\}\Psi = 0 \quad (1.1)$$

to a new equivalent equation according the following scheme

$$\{D\}\Psi = 0 \Rightarrow \mathcal{F}\{D\}\Gamma\Gamma^{-1}\Psi = (\hat{K}_1 + \hat{K}_2)\Phi = 0 \quad (1.2)$$

with

$$\Psi = \Gamma\Phi \quad (1.3)$$

where $\hat{K}_1$ and $\hat{K}_2$ are first-order commuting matrix differential operators

$$[\hat{K}_1, \hat{K}_2]_- = 0 \quad (1.4)$$

depending each one on different space-time variables, Γ is a constant nonsingular matrix, and $\mathcal{F}$ that we have to find (if they exist) in order to fulfill the condition (1.4). Depending on the problem to solve there are two different schemes of separation of variables (1) Consecutive separation of variables when one variable is separated from the remaining three variables, then one variable is separated from the other two variables, and finally, the remaining two variables are separated. (2) Pairwise separation of variables, when two space-time variables are separated from the another two ones, and the resulting operators are also separated. In the present article we focus our attention to the Dirac equation in the presence of a central field, like the Hydrogen Atom where a complete separation of variables is possible. In this case, using a pairwise scheme of separation, the angular dependence is completely separated from the other two variables. The article is structured as follows. In Sec. II we separate variables in the Dirac equation. In Sec III. we solve the angular equation in the local rotating frame in terms of Jacobi Polynomials In Sec. IV we establish the relation between the spinor solution in the diagonal (rotating) tetrad gauge and the fixed Cartesian one.

2 Separation of variables

In this section we proceed to separate variables in the Dirac equation (1.1), where $\tilde{\gamma}^\mu$ are the curved gamma matrices satisfying the relation,

$$\{\tilde{\gamma}^\mu, \tilde{\gamma}^\nu\}_+ = 2g^{\mu\nu} \quad (2.1)$$

and Γ_μ are the spin connections [7].

$$\Gamma_\lambda = \frac{1}{4}g_{\mu\alpha}[(\partial b_\nu^\beta/\partial x^\lambda)a_\beta^\alpha - \Gamma_{\nu\lambda}^\alpha]s^{\mu\nu} \quad (2.2)$$

with

$$s^{\mu\nu} = \frac{1}{2}(\tilde{\gamma}^\mu\tilde{\gamma}^\nu - \tilde{\gamma}^\nu\tilde{\gamma}^\mu) \quad (2.3)$$

and the matrices b_α^β , a_β^α establish the connection between the Dirac matrices ($\tilde{\gamma}$) on a curved spacetime and the Minkowski space (γ) Dirac matrices as follows:

$$\tilde{\gamma}_\mu = b_\mu^\alpha \gamma_\alpha \quad \tilde{\gamma}^\mu = a_\beta^\mu \gamma^\beta \quad (2.4)$$

If we choose to work in the fixed Cartesian gauge, then spinor connections are zero and the $\tilde{\gamma}$ matrices take the form

$$\begin{aligned} \tilde{\gamma}^0 &= \gamma^0 = \bar{\gamma}^0, \quad \tilde{\gamma}^1 = \left[(\gamma^1 \cos \varphi + \gamma^2 \sin \varphi) \sin \vartheta + \gamma^3 \cos \vartheta \right] = \bar{\gamma}^1, \\ \tilde{\gamma}^2 &= \frac{1}{r} \left[(\gamma^1 \cos \varphi + \gamma^2 \sin \varphi) \cos \vartheta - \gamma^3 \sin \vartheta \right] = \frac{\bar{\gamma}^2}{r}, \\ \tilde{\gamma}^3 &= \frac{1}{r \sin \vartheta} (-\gamma^1 \sin \varphi + \gamma^2 \cos \varphi) = \frac{\bar{\gamma}^3}{r \sin \vartheta} \end{aligned} \quad (2.5)$$

and the Dirac equation in the fixed tetrad frame (2.5) takes the form

$$\left\{ \bar{\gamma}^0 (\partial_t + iV(r)) + \bar{\gamma}^1 \partial_r + \frac{\bar{\gamma}^2}{r} \partial_\vartheta + \frac{\bar{\gamma}^3}{r \sin \vartheta} \partial_\varphi + m \right\} \tilde{\Psi} = 0 \quad (2.6)$$

In order to separate variables in the Dirac equation, we are going to work in the diagonal tetrad gauge where the gamma matrices $\tilde{\gamma}_d$ take the form

$$\tilde{\gamma}_d^0 = \gamma^0, \quad \tilde{\gamma}_d^1 = \gamma^1, \quad \tilde{\gamma}_d^2 = \frac{1}{r} \gamma^2, \quad \tilde{\gamma}_d^3 = \frac{1}{r \sin \vartheta} \gamma^3 \quad (2.7)$$

Since the curvilinear matrices $\tilde{\gamma}^\mu$ matrices and $\tilde{\gamma}_d$ satisfy the same anticommutation relations, they are related by a similarity transformation, unique up to a factor. In the present case we choose this factor in order to eliminate the spin connections in the resulting Dirac equation. The transformation S can be written as [8]

$$S = \frac{1}{r(\sin \vartheta)^{1/2}} \exp\left(-\frac{\varphi}{2} \gamma^1 \gamma^2\right) \exp\left(-\frac{\vartheta}{2} \gamma^3 \gamma^1\right) \mathbf{a} = S_0 \mathbf{a} \quad (2.8)$$

where $\mathbf{a}$ is the constant non singular matrix given by the expression

$$\mathbf{a} = \frac{1}{2} (\gamma^1 \gamma^2 - \gamma^1 \gamma^3 + \gamma^2 \gamma^3 + I) \quad (2.9)$$

which applied on the gamma's acts as follows

$$\mathbf{a} \gamma^1 \mathbf{a}^{-1} = \gamma^3, \quad \mathbf{a} \gamma^2 \mathbf{a}^{-1} = \gamma^1, \quad \mathbf{a} \gamma^3 \mathbf{a}^{-1} = \gamma^2, \quad (2.10)$$

the transformation S acts on the curvilinear $\tilde{\gamma}$ matrices, reducing them to the rotating diagonal gauge as follows

$$S^{-1}\tilde{\gamma}^\mu S = g^{\mu\mu}\gamma^\mu = \tilde{\gamma}_d^\mu \quad (\text{no summation}) \quad (2.11)$$

then, the Dirac equation in spherical coordinates, with the radial potential $V(r)$, in the local rotating frame reads

$$\left\{ \gamma^0 \partial_t + \gamma^1 \partial_r + \frac{\gamma^2}{r} \partial_\vartheta + \frac{\gamma^3}{r \sin \vartheta} \partial_\varphi + m + i\gamma^0 V(r) \right\} \Psi = 0 \quad (2.12)$$

where we have introduced the spinor Ψ , related to $\tilde{\Psi}$ by the expression

$$\tilde{\Psi} = S\Psi = S_0 \mathbf{a} \Psi \quad (2.13)$$

and γ^μ are the standard Dirac flat matrices. Applying the algebraic method of separation of variables [9, 10], it is possible to write eq. (2.12) as a sum of two first order linear differential operators $\hat{K}_1$, $\hat{K}_2$ satisfying the relation

$$[\hat{K}_1, \hat{K}_2] = 0, \quad \{ \hat{K}_1 + \hat{K}_2 \} \Phi = 0 \quad (2.14)$$

$$\hat{K}_1 \Phi = \lambda \Phi = -\hat{K}_2 \Phi \quad (2.15)$$

then, if we separate the time and radial dependence from the angular one, we obtain

$$\hat{K}_2 \Phi = \left[\gamma^2 \partial_\vartheta + \frac{\gamma^3}{\sin \vartheta} \partial_\varphi \right] \gamma^0 \gamma^1 \Phi = -i\kappa \Phi \quad (2.16)$$

$$\hat{K}_1 \Phi = r \left[\gamma^0 \partial_t + \gamma^1 \partial_r + m + i\gamma^0 V \right] \gamma^0 \gamma^1 \Phi = i\kappa \Phi \quad (2.17)$$

with

$$\Psi = \gamma^0 \gamma^1 \Phi \quad (2.18)$$

where we have made the identification $i\kappa = \lambda$. Notice that (2.16) is the angular momentum $\hat{K}$ obtained by Brill and Wheeler [7]. Here it is necessary to remark that the operator $\hat{K}_2$ appearing in (2.16) is not singled valued and does not satisfy the properties of a "good"

angular momentum operator. The true angular operator should be obtained from $\hat{K}_2$ with the help of the transformation S (2.8)

$$S \left[\gamma^2 \partial_\vartheta + \frac{\gamma^3}{\sin \vartheta} \partial_\varphi \right] \gamma^0 \gamma^1 S^{-1} S \Phi = -i\kappa S \Phi \quad (2.19)$$

since $S\gamma^\mu S^{-1} = \bar{\gamma}^\mu$, we obtain

$$\left[\bar{\gamma}^2 \bar{\gamma}^0 \bar{\gamma}^1 \partial_\vartheta + \frac{\bar{\gamma}^3 \bar{\gamma}^0 \bar{\gamma}^1}{\sin \vartheta} \partial_\varphi + \bar{\gamma}^2 \bar{\gamma}^0 \bar{\gamma}^1 S \partial_\vartheta S^{-1} + \frac{\bar{\gamma}^3 \bar{\gamma}^0 \bar{\gamma}^1}{\sin \vartheta} S \partial_\varphi S^{-1} \right] \tilde{\Phi} = -i\kappa \tilde{\Phi} \quad (2.20)$$

where $S\Phi = \tilde{\Phi}$ using the explicit form of the transformation S given by (2.8) we have

$$S \partial_\vartheta S^{-1} = \frac{1}{2} \cot \vartheta + \frac{1}{2} (\cos \varphi - \gamma^1 \gamma^2 \sin \varphi) \gamma^3 \gamma^1 \quad (2.21)$$

$$S \partial_\varphi S^{-1} = \frac{1}{2} \gamma^1 \gamma^2 \quad (2.22)$$

substituting (2.21) and (2.22) into (2.20)

$$\left[\bar{\gamma}^0 \bar{\gamma}^2 \bar{\gamma}^1 \partial_\vartheta + \frac{\bar{\gamma}^0 \bar{\gamma}^3 \bar{\gamma}^1}{\sin \vartheta} \partial_\varphi + \bar{\gamma}^0 \right] \tilde{\Phi} = i\kappa \tilde{\Phi} \quad (2.23)$$

expression that can be written as follows

$$\hat{K} = \gamma^0 [(\sigma \hat{\mathbf{L}}) + 1] \quad (2.24)$$

The operator $\hat{K}$ satisfies the relation

$$\hat{K}^2 = \hat{J}^2 + \frac{1}{4}, \quad \hat{J}^2 = (L + \frac{1}{2}\sigma)^2 \quad (2.25)$$

and the eigenvalue k is related to j as follows

$$k^2 = j(j+1) + \frac{1}{2} = (j + \frac{1}{2})^2 \quad (2.26)$$

Now we proceed to decouple the equation (2.16) governing the angular dependence of the Dirac spinor. In order to simplify the resulting equations, we choose to work with the auxiliary spinor $\bar{\Phi}$ related to Φ as follows

$$\bar{\Phi} = \mathbf{a} \Phi \quad (2.27)$$

then the equation (2.16) takes the form

$$\left[\gamma^1 \partial_\vartheta + \frac{\gamma^2}{\sin \vartheta} \partial_\varphi \right] \gamma^0 \gamma^3 \bar{\Phi} = -i\kappa \bar{\Phi} \quad (2.28)$$

In order to reduce the equation (2.28) to a system of ordinary differential equations, we choose to work in the following representation for the gamma matrices, [11]

$$\gamma^i = \begin{pmatrix} 0 & \sigma^i \\ \sigma^i & 0 \end{pmatrix}, \gamma^0 = \begin{pmatrix} -i & 0 \\ 0 & i \end{pmatrix}, \quad (2.29)$$

Then, substituting (2.29) into (2.28) we obtain,

$$\left[\sigma^2 \partial_\vartheta - i \frac{\sigma^1}{\sin \vartheta} m \right] \bar{\Phi}_1 = -ik \bar{\Phi}_1 \quad (2.30)$$

$$\left[-\sigma^2 \partial_\vartheta + i \frac{\sigma^1}{\sin \vartheta} m \right] \bar{\Phi}_2 = -ik \bar{\Phi}_2 \quad (2.31)$$

with

$$\bar{\Phi} = \begin{pmatrix} \bar{\Phi}_1 \\ \bar{\Phi}_2 \end{pmatrix} \quad (2.32)$$

3 Solution of the angular equations

In this section we are going to solve the systems of equations (2.30) and (2.31). It is not difficult to see that the spinor $\bar{\Phi}_2$ is proportional to $\sigma^3 \bar{\Phi}_1$, this property allows us to consider only the system (2.30). Using the standard Pauli matrices, we have that (2.30) reduces to

$$k \sin \vartheta \xi_1 - m \xi_2 - \sin \vartheta \frac{d\xi_2}{d\vartheta} = 0 \quad (3.1)$$

$$k \sin \vartheta \xi_2 - m \xi_1 + \sin \vartheta \frac{d\xi_1}{d\vartheta} = 0 \quad (3.2)$$

with

$$\bar{\Phi}_1 = \begin{pmatrix} \xi_1 \\ \xi_2 \end{pmatrix} \quad (3.3)$$

In order to solve the coupled system of equations (3.1)-(3.2) we make the following ansatz,

$$\xi_1 = i(\sin \frac{\vartheta}{2})^m (\cos \frac{\vartheta}{2})^{m+1} q(\vartheta), \quad \xi_2 = (\sin \frac{\vartheta}{2})^{m+1} (\cos \frac{\vartheta}{2})^m f(\vartheta) \quad (3.4)$$

Substituting (3.4) into (3.1) and (3.2) we obtain,

$$ikq(x) - (m + \frac{1}{2})f(x) + (1-x)\frac{df(x)}{dx} = 0 \quad (3.5)$$

$$ikf(x) + (m + \frac{1}{2})q(x) + (1+x)\frac{dq(x)}{dx} = 0 \quad (3.6)$$

where we have made the change of variable $x = \cos \vartheta$, The normalizable solution of the system (3.5)-(3.6) can be expressed in terms of Jacobi Polynomials $P_k^{(\alpha, \beta)}$ [12]

$$\xi_1 = c_0 (\sin \frac{\vartheta}{2})^m (\cos \frac{\vartheta}{2})^{m+1} P_n^{(m-1/2, m+1/2)} \quad (3.7)$$

$$\xi_2 = c_0 (\sin \frac{\vartheta}{2})^{m+1} (\cos \frac{\vartheta}{2})^m P_n^{(m+1/2, m-1/2)} \quad (3.8)$$

where c_0 is an arbitrary constant, which can be fixed in order to normalize the angular dependence of the spinor Φ according to the product

$$2\pi \int_0^{2\pi} \Phi_k^{\mu\dagger} \Phi_{k'}^\mu d\vartheta = \delta_{kk'} \quad (3.9)$$

and n is given by the expression

$$n = -m - \frac{1}{2} + k, \quad n = 0, 1, 2, \dots \quad (3.10)$$

then, using the orthogonality relation for the Jacobi polynomials [12]

$$\begin{aligned} \int_{-1}^1 (1-x)^\alpha (1+x)^\beta P_n^{(\alpha, \beta)}(x) P_m^{(\alpha, \beta)}(x) dx = \\ \frac{2^{\alpha+\beta+1}}{2n+\alpha+\beta+1} \frac{\Gamma(n+\alpha+1)\Gamma(n+\beta+1)}{\Gamma(n+1)\Gamma(n+\alpha+\beta+1)} \delta_{mn} \end{aligned} \quad (3.11)$$

we obtain

$$c_0 = \frac{\sqrt{n!(n+2m)!}}{(n+m-1/2)!\sqrt{2\pi}} \quad (3.12)$$

Now, we are going to solve the system of equation (2.17), governing the radial dependence of the spinor Ψ solution of the Dirac equation. This equation can be written in the form,

$$\left(-\gamma^3\partial_t + \gamma^0\partial_r + m\gamma^0\gamma^3 - i\gamma^3V(r) + i\frac{\kappa}{r}\right)\bar{\Phi} = 0 \quad (3.13)$$

Using the representation for the gamma matrices given by (2.29), we obtain the following system of equations

$$\left(d_r + \frac{k}{r}\right)\bar{\Phi}_1 - \sigma^3(E - V(r) - m)\bar{\Phi}_2 = 0 \quad (3.14)$$

$$\left(-d_r + \frac{k}{r}\right)\bar{\Phi}_2 - \sigma^3(E - V(r) + m)\bar{\Phi}_1 = 0 \quad (3.15)$$

therefore, the general form of the spinor $\bar{\Phi}$ is given by the expression

$$\bar{\Phi} = c_0 \left(\sin \frac{\vartheta}{2}\right)^m \left(\cos \frac{\vartheta}{2}\right)^m e^{i(m\varphi - Et)} \begin{pmatrix} c(\cos \frac{\vartheta}{2})P_n^{(m-1/2, m+1/2)}A(r) \\ c(\sin \frac{\vartheta}{2})P_n^{(m+1/2, m-1/2)}A(r) \\ (\cos \frac{\vartheta}{2})P_n^{(m-1/2, m+1/2)}B(r) \\ -(\sin \frac{\vartheta}{2})P_n^{(m+1/2, m-1/2)}B(r) \end{pmatrix} \quad (3.16)$$

where c_0 is given by (3.12), c is a constant, and $A(r), B(r)$ satisfy the system of equations

$$\left(d_r + \frac{k}{r}\right)A(r) - (E - V(r) - m)B(r) = 0 \quad (3.17)$$

$$\left(-d_r + \frac{k}{r}\right)B(r) - (E - V(r) + m)A(r) = 0 \quad (3.18)$$

obviously, the form $A(r)$ and $B(r)$ will depend on the potential $V(r)$. Among the solvable problems we have the Coulomb case where the radial dependence can be expressed in terms of confluent hypergeometric functions.[13]

Regarding the eigenvalues m of the projection of the angular momentum operator $-i\partial_\varphi$ we have that since

$$S_z(\varphi + \pi) = -S_z(\varphi) \quad (3.19)$$

and the spinor $\tilde{\Psi}$ is single valued, then $\Psi(\varphi + 2\pi) = -\Psi(\varphi)$, and therefore m takes half integer values

$$m = N + \frac{1}{2}, \quad N = 0, \pm 1, \pm 2 \dots \quad (3.20)$$

4 Transformation of the spinor

In this section we proceed to show that the angular dependence of the spinor solution of the Dirac equation (2.12), which is given by the expression (3.16), reduces to the standard spherical spinors when we apply the transformation S given by (2.8). Since the spinor $\tilde{\Psi}$, expressed in the Cartesian tetrad gauge, is related to the spinor solution Ψ in the local rotating frame by means of (2.13), we have

$$S_0 \gamma^0 \gamma^3 \bar{\Phi} = \tilde{\Psi} \quad (4.1)$$

where we have used (2.18) and (2.10). Using the representation (2.29) and the spinor (3.16), we obtain

$$\tilde{\Psi} = \begin{pmatrix} \tilde{\Psi}^+ \\ \tilde{\Psi}^- \end{pmatrix} - i \frac{c_0}{\sqrt{2}r} \left(\sin \frac{\vartheta}{2} \cos \frac{\vartheta}{2} \right)^{m-1/2} S_{\vartheta}(\vartheta) e^{im\varphi} \begin{pmatrix} (\cos \frac{\vartheta}{2}) P_n^{(m-1/2, m+1/2)} B(r) \\ (\sin \frac{\vartheta}{2}) P_n^{(m+1/2, m-1/2)} B(r) \\ -c(\cos \frac{\vartheta}{2}) P_n^{(m-1/2, m+1/2)} A(r) \\ c(\sin \frac{\vartheta}{2}) P_n^{(m+1/2, m-1/2)} A(r) \end{pmatrix} \quad (4.2)$$

$$S_{\varphi}(\varphi) S_{\vartheta}(\vartheta) = \begin{pmatrix} e^{-i\varphi/2} & 0 & 0 & 0 \\ 0 & e^{i\varphi/2} & 0 & 0 \\ 0 & 0 & e^{-i\varphi/2} & 0 \\ 0 & 0 & 0 & e^{i\varphi/2} \end{pmatrix} \begin{pmatrix} \cos \frac{\vartheta}{2} & \sin \frac{\vartheta}{2} & 0 & 0 \\ -\sin \frac{\vartheta}{2} & \cos \frac{\vartheta}{2} & 0 & 0 \\ 0 & 0 & \cos \frac{\vartheta}{2} & \sin \frac{\vartheta}{2} \\ 0 & 0 & -\sin \frac{\vartheta}{2} & \cos \frac{\vartheta}{2} \end{pmatrix} \quad (4.3)$$

substituting (4.3) into (4.2) and taking into account the relations

$$(1-x)P_n^{(\alpha+1, \beta)}(x) + (1+x)P_n^{(\alpha, \beta+1)}(x) = 2P_n^{(\alpha, \beta)}(x) \quad (4.4)$$

$$P_n^{(\alpha, \beta-1)}(x) = P_n^{(\alpha-1, \beta)}(x) + P_{n+1}^{(\alpha, \beta)}(x) \quad (4.5)$$

we find that the upper two components of the spinor $\tilde{\Psi}$ (4.2) take the form

$$\tilde{\Psi}^+ = -i \frac{c_0}{\sqrt{2}r} \left(\sin \frac{\vartheta}{2} \cos \frac{\vartheta}{2} \right)^{1/2} \begin{pmatrix} \left(\sin \frac{\vartheta}{2} \cos \frac{\vartheta}{2} \right)^{m-1/2} P_n^{(m-1/2, m-1/2)}(\cos \vartheta) e^{i(m-1/2)\varphi} B(r) \\ \left(\sin \frac{\vartheta}{2} \cos \frac{\vartheta}{2} \right)^{m+1/2} P_{n-1}^{(m+1/2, m+1/2)}(\cos \vartheta) e^{i(m+1/2)\varphi} B(r) \end{pmatrix} \quad (4.6)$$

with the help of the relation

$$Y_{l,m}(\vartheta, \varphi) = \frac{1}{2^m l!} \sqrt{\frac{2l+1}{4\pi}} (l-m)!(l+m)!(1-x^2)^{m/2} P_{l-m}^{(m,m)}(x) e^{im\varphi} \quad (4.7)$$

and

$$P_l^{-m}(x) = (-1)^m \frac{(l-m)!}{(l+m)!} P_l^m(x) \quad (4.8)$$

we find that $\tilde{\Psi}^+$ can be expressed in terms of spherical harmonics $Y_{k,m}(\vartheta, \varphi)$ as follows

$$\tilde{\Psi}^+ = -i \frac{1}{r} (2k+1)^{-1/2} \begin{pmatrix} (k+m+1/2)^{1/2} Y_k^{m-1/2}(\vartheta, \varphi) B(r) \\ (k-m+1/2) Y_k^{m+1/2}(\vartheta, \varphi) B(r) \end{pmatrix} \quad (4.9)$$

since the transformation S_0 acting on the spinor Ψ has a block form, it is possible to write it as follows

$$S_0 = \begin{pmatrix} \hat{S}_0 & \\ & \hat{S}_0 \end{pmatrix}, \quad \hat{S}_0 = \frac{1}{r(\sin \vartheta)^{1/2}} \begin{pmatrix} e^{-i\varphi/2} & 0 \\ 0 & e^{i\varphi/2} \end{pmatrix} \begin{pmatrix} \cos \frac{\vartheta}{2} & \sin \frac{\vartheta}{2} \\ -\sin \frac{\vartheta}{2} & \cos \frac{\vartheta}{2} \end{pmatrix} \quad (4.10)$$

then, if we set $c=-1$, we have

$$\tilde{\Psi}^- = \hat{S}_0 \bar{\Phi}^- = \hat{S}_0 \sigma_3 \bar{\Phi}^+ = \hat{S}_0 \sigma^3 \hat{S}_0^{-1} \hat{S}_0 \bar{\Phi}^+ = \sigma \cdot \hat{\mathbf{e}}_r \tilde{\Psi}^+ \quad (4.11)$$

and the lower components $\tilde{\Psi}^-$ of the spinor $\tilde{\Psi}$ read

$$\tilde{\Psi}^- = i \frac{1}{r} (2k+1)^{-1/2} \begin{pmatrix} (k-m+1/2)^{1/2} Y_k^{m-1/2}(\vartheta, \varphi) A(r) \\ -(k+m+1/2) Y_k^{m+1/2}(\vartheta, \varphi) A(r) \end{pmatrix} \quad (4.12)$$

the results (4.9) and (4.12) are in terms of the well known spherical spinors as in almost all the textbooks on relativistic quantum mechanics. The basis solution in terms of Jacobi Polynomials has been of utility in analyzing the Dirac spinor in the presence of a magnetic monopole, also it can be of help in discussing non-central interactions.

acknowledgments

The author wishes to express his indebtedness to the Centre de Physique Théorique for the suitable conditions of work. Also the author wishes to acknowledge to the CONICIT of Venezuela and to the Fundación Polar for financial support.

References

- [1] W. Gordon, “Die Energienveans des Wasserstoffatoms nach der Diracschen Quantentheorie des Electrons” Z. Phys. **48** 11-14 (1928)
- [2] D. V. Volkov, “Über eine Klasse von Losungen der Diracschen Gleichung” Z. Phys. **94**, 3-4 (1935)
- [3] V. G. Bagrov, and D. M. Gitman *Exact Solutions of Relativistic Wave Equations* (Kluwer, Dordretch, 1990)
- [4] W. Miller Jr. In *Symmetries and nonlinear phenomena* (World Scientific, Singapore 1988)
- [5] W. Miller Jr. *Symmetry and separation of variables* (Addisson Wesley, Massachusetts, 1977)
- [6] A. H. Cook, “On separable solutions of Dirac’s equation for the electron” Proc. R. Soc. A. **383** 247-278 (1982)
- [7] D. Brill and J. A. Wheeler, “Interaction of Neutrinos and Gravitational Fields” Rev. Mod. Phys. **29**, 465-479 (1957)
- [8] G. V. Shishkin, V. M. Villalba, “Separation of variables of the Dirac equation in the presence of external scalar fields” Vestn. Beloruss. Gos. Univ. **3** 6-12 (1987)
- [9] G.V. Shishkin and V.M. Villalba, “Dirac equation in external vector fields: separation of variables” J. Math. Phys. **30**, 2132-2143 (1989)
- [10] G.V. Shishkin and V.M. Villalba, “Neutrino in the presence of gravitational fields: Separation of variables” J. Math. Phys. **33**, 2093-2110 (1992)

- [11] J.M. Jauch and F. Rohrlich, *The Theory of Photons and Electrons* (Springer, New York, 1976)
- [12] W. Magnus, F. Oberhettinger, and R. P. Soni *Formulas and Theorems for the Special Functions of Mathematical Physics* (Springer-Verlag, Berlin, 1966)
- [13] A. S. Davydov, *Quantum Mechanics* (Pergamon, Oxford 1965)